



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE. I was not aware of the utility, until I saw Mr. Evans' rule, of any rule for the extraction of roots above the common one. I have derived his formula, and also the following which will give the value of R to one-half more decimal places than Mr. Evans' rule:

$$R = r \times \frac{(n+1)N + (n-1)r^n}{(n-1)N + (n+1)r^n},$$

where the letters represent the same numbers as in the formula given by Mr. Evans, ANALYST, page 11, Vol. III.

Example. To find the square root of 2. Here $n=2$, $N=2$, and let the approximate trial root be 1.4, then

$$R = 1.4 \times \frac{3 \times 2 + (1.4)^2}{2 + 3(1.4)^2} = 1.4 \times \frac{7.96}{7.88} = 1.4 \times \frac{199}{197} = 1.4142133$$

correct to 6 places of decimals. In this formula, if r be too great R will also, and vice versa, and therefore by using two trial roots, one too great and the other too small, the limits, between which the true root is, are obtained.

R. J. Adcock.

[98. Mr. Adcock objects to the published solution of 98. He finds the height due the velocity of the stream (13.2 feet per sec.) to be 2.7+ feet. And because the pressure of a fluid column of that height would produce, in an equal sectional area of the stream, the given velocity, he contends that this pressure, viz., $2.7+ \times 62\frac{1}{2} = 169.2$ is the correct answer.]

BOOK NOTICE.

Elements of Geometry With Exercises for Students, and an Introduction to Modern Geometry, By A. SCHUYLER, LL. D., President of Baldwin University, &c.

This book will interest students on account of its *superior* mechanical execution; but it is especially valuable for the logical accuracy with which the various propositions are announced and demonstrated. The Introduction to Modern Geometry is, for many students, a valuable addition.

ERRATA.

On page 35, erase lines 9, 10 and 11.

“ “ 41, “ third member in lines 9, 10 and 11.

“ “ 53, lines 18, 20 and 22, for $X(a_1$ or $b_1)$ read $X_1(a_1$ or $b_1)$.

“ “ 63, last line, for constant angle at N , &c., read constant length of NC .

“ “ 75, line 5 from bottom for $\frac{ec}{bh}e$, read $\frac{ec}{bh}a$.